

An optimized chiral nucleon-nucleon interaction at next-to-next-to-leading order

A. Ekström,^{1,2} G. Baardsen,³ C. Forssén,³ G. Hagen,^{4,5} M. Hjorth-Jensen,⁶ G. R. Jansen,⁷
R. Machleidt,⁷ W. Nazarewicz,⁸ T. Papenbrock,⁹ J. Sarich,⁹ and S. M. Wild⁹

¹*Department of Physics and Center of Mathematics for Applications, University of Oslo, N-0316 Oslo, Norway*

²*National Superconducting Cyclotron Laboratory, Michigan State University, East Lansing, MI 48824-1321, USA*

³*Department of Fundamental Physics, Chalmers University of Technology, SE-412 96 Göteborg, Sweden*

⁴*Physics Division, Oak Ridge National Laboratory, Oak Ridge, TN 37831, USA*

⁵*Department of Physics and Astronomy, University of Tennessee, Knoxville, TN 37996, USA*

⁶*Department of Physics and Astronomy, Michigan State University, East Lansing, MI 48824, USA*

⁷*Department of Physics, University of Idaho, Moscow, ID 83844, USA*

⁸*Faculty of Physics, University of Warsaw, ul. Hoża 69, 00-681 Warsaw, Poland*

⁹*Mathematics and Computer Science Division, Argonne National Laboratory, Argonne, IL 60439, USA*

We optimize the nucleon-nucleon interaction from chiral effective field theory at next-to-next-to-leading order. The resulting new chiral force NNLO_{opt} yields $\chi^2 \approx 1$ per degree of freedom for laboratory energies below approximately 125 MeV. In the $A = 3, 4$ nucleon systems, the contributions of three-nucleon forces are smaller than for previous parametrizations of chiral interactions. We use NNLO_{opt} to study properties of key nuclei and neutron matter, and demonstrate that many aspects of nuclear structure can be understood in terms of this nucleon-nucleon interaction, without explicitly invoking three-nucleon forces.

PACS numbers: 21.30.-x, 21.10.-k, 21.45.-v, 21.60.-n

Introduction. – Interactions from chiral effective field theory (EFT) employ symmetries and the pattern of spontaneous symmetry breaking of quantum chromodynamics [1, 2]. In this approach, the exchange of pions within chiral perturbation theory yields the long-ranged contributions of the nuclear interaction, while short-ranged components are included as contact terms. The interaction is parametrized in terms of low-energy constants (LECs) that are determined by fit to experimental data. The interactions from chiral EFT exhibit a power counting in the ratio Q/Λ , with Q being the low-momentum scale being probed and Λ the cutoff which is of the order of 1 GeV. At next-to-next-to-leading order (NNLO), three-nucleon forces (3NFs) enter, while four-nucleon forces (4NFs) enter at next-to-next-to-next-to-leading order (N³LO). For laboratory energies below 125 MeV, the nucleon-nucleon (NN) force exhibits a quality of fit with $\chi^2 \approx 10/\text{datum}$ at NNLO [3], while a high-precision potential N³LO_{EM}, with a $\chi^2 \approx 1/\text{datum}$ up to 290 MeV, was obtained by Entem and Machleidt [2, 4].

The 3NFs at NNLO that accompany the current N³LO NN potentials play a pivotal role in nuclear structure calculations [5]. They determine the ground-state spin of ¹⁰B [6], correctly set the drip line in oxygen isotopes [7, 8], and make ⁴⁸Ca a doubly magic nucleus [9, 10]. While it might seem surprising that smaller corrections at NNLO are so decisive for basic nuclear structure properties, the 3NF contains spin-orbit and tensor contributions that clearly are important for the currently employed chiral interactions. The contributions of 3NFs at N³LO have also been worked out [11, 12], and there are on-going efforts to compute even higher orders [13].

While the quest for higher orders is important, this approach will only result in higher accuracy if the optimization at lower orders were carried out accurately. This makes it important and timely to revisit the optimization question. We note in particular that the fits of the currently employed chiral interactions [3, 4, 14] date back about a decade, and that there has been a considerable recent progress in developing tools for the derivative-free nonlinear least-squares optimization [15]. Furthermore, the quantification of theoretical uncertainties is a long-term objective of nuclear structure theory, and this requires a covariance analysis of the interaction parameters with respect to the experimental uncertainties of the nucleon-nucleon elastic scattering observables, see, e.g., Refs. [15, 16]. It is the purpose of this Letter to take the first step toward this goal. In what follows, we present a state-of-the-art optimization of the NN chiral EFT interaction at NNLO. This yields a much-improved χ^2 and a high-precision NN potential NNLO_{opt}. The 3NF at NNLO is adjusted to the binding energies in $A = 3, 4$ nuclei. We present computations of three-nucleon and four-nucleon bound states, and employ NNLO_{opt} to ground- and excited states in ¹⁰B, masses and excited states of oxygen and calcium isotopes, and neutron matter.

Optimizing the NN interaction at NNLO. – For the optimization of the chiral NN interaction we use the Practical Optimization Using No Derivatives (for Squares) algorithm POUNDerS [15] as implemented in [17]. This derivative-free algorithm employs a quadratic model and is particularly useful for computationally expensive objective functions. We optimize the three pion-nucleon (πN) couplings (c_1, c_3, c_4), and 11 partial wave contact parameters C and \bar{C} , while we keep the axial-vector cou-

pling constant g_A , the pion-decay constant f_π , and all masses fixed. In the optimization, we minimize the objective function

$$f(\vec{x}) = \sum_{q=1}^{N_q} \left(\frac{\delta_q^{\text{NNLO}}(\vec{x}) - \delta_q^{\text{Nijm93}}}{w_q} \right)^2, \quad (1)$$

where δ^{NNLO} are NNLO phase shifts, δ^{Nijm93} are experimental phase shifts from the Nijmegen multi-energy partial-wave analysis [18], \vec{x} denotes the parameters of the chiral interaction, and w_q are weighting factors. Note that Eq. (1) is not the χ^2 with respect to experimental data. The actual χ^2 is calculated following the POUNDerS optimization. The phase shifts δ^{NNLO} are computed from R -matrix inversion, and in the proton-proton (pp) channels we include the Coulomb interaction [19, 20]. The contact terms are optimized to reproduce the Nijmegen phase shifts for each corresponding partial wave, while keeping the c_i 's fixed. For the contacts, the weight w_q scales with the third power of the relative momentum q , while for the c_i 's, we employ the uncertainties quoted in the Nijmegen analysis [18]. This can be justified by a physical argument: for the peripheral waves the higher energies still represent longer-range physics, and the need for a pedantic agreement with lower energy phase shifts can be weakened. The πN couplings c_1 , c_3 , and c_4 were simultaneously optimized to the peripheral partial-waves 1D_2 , 3D_2 , 3F_2 , E_2 , 3F_3 , 1G_4 , and 3F_4 . Note that the NNLO contact terms do not contribute for orbital angular momenta $L \geq 2$. We do not include other peripheral waves from the Nijmegen study since they carry extremely small uncertainties, which leads to a very noisy objective function.

Table I summarizes the optimization results. Our values should be compared to the πN couplings as determined from πN scattering data, where $c_1 = -0.81 \pm 0.15$, $c_3 = -4.69 \pm 1.34$, and $c_4 = +3.40 \pm 0.04$ has been obtained [21]. Thus, POUNDerS yields values for c_1 and c_3 which agree well with the empirical determination from πN scattering. The c_4 value, however, deviates significantly from its empirical value. The same trend was also found in the construction of the $N^3\text{LO}$ [4] NN interaction. A detailed statistical sensitivity analysis of the LECs with uncertainty quantification will be presented in Ref. [22].

Table II shows the χ^2/datum for NNLO_{opt} at various laboratory energy bins. The quality of the fit is particularly good for energies below 125 MeV. For comparison, the np NNLO interaction of Ref. [3] yields χ^2/datum of 12–27 in the range $\Lambda = 600/700 - 450/500$ MeV at energies up to 290 MeV.

Around energies of 144 MeV there exist two data sets of pp differential cross sections with a very high precision (0.5% error) [24] (47 data points). The total number of pp data in the energy interval 125–183 MeV is 343. The unusual precision of those 47 data points distorts

TABLE I. Parameters of NNLO_{opt} at $\Lambda = 500$ MeV: c_i (in GeV^{-1}), \tilde{C} (in 10^4GeV^{-2}), and C (in 10^4GeV^{-4}).

LEC	value	LEC	value	LEC	value
c_1	-0.91863953	c_3	-3.88868749	c_4	4.31032716
$\tilde{C}_{1S_0}^{pp}$	-0.15136604	$\tilde{C}_{1S_0}^{np}$	-0.15214109	$\tilde{C}_{1S_0}^{nn}$	-0.15176475
C_{1S_0}	2.40402194	C_{3S_1}	0.92838466	\tilde{C}_{3S_1}	-0.15843418
C_{1P_1}	0.41704554	C_{3P_0}	1.26339076	C_{3P_1}	-0.78265850
$C_{3S_1-3D_1}$	0.61814142	C_{3P_2}	-0.67780851		

TABLE II. χ^2/datum for NNLO_{opt} at $\Lambda = 500$ MeV with respect to the np and pp 1999 databases [23]. The values without the high precision data sets [24] are marked by asterisks.

T_{lab} (MeV)	0–35	35–125	125–183	183–290	0–290
pp χ^2/datum	1.11	1.56	$\left\{ \begin{array}{l} 23.95 \\ 4.35^* \end{array} \right.$	29.26	$\left\{ \begin{array}{l} \mathbf{17.10} \\ \mathbf{14.03}^* \end{array} \right.$
np χ^2/datum	0.85	1.17	1.87	6.09	2.95

the χ^2/datum for this interval. For this reason, Table II also shows the results without the high-precision data.

Two comments are in order. First, the χ^2 with respect to scattering observables is lower when the 1P_1 phase shifts are weighted with the uncertainties from the Nijmegen analysis. The P -waves are only accurately reproduced when going to $N^3\text{LO}$ [4]. Second, the $^3S_1 - ^3D_1$ coupled channel is optimized with the additional constraint of reproducing the deuteron binding energy. The remaining deuteron observables, as well as the 1S_0 scattering observables are predictions and reproduce the experimental values well, see Table III.

TABLE III. Scattering lengths a and effective ranges r (both in fm). The superscripts N and C for the proton-proton observables refer to nuclear forces and Coulomb-plus-nuclear forces, respectively. B_D , r_D , Q_D , and P_D denote the deuteron binding energy, radius, quadrupole moment, and D -state probability, respectively. Q_D and r_D are calculated without meson-exchange currents and relativistic corrections.

	$N^3\text{LO}_{\text{EM}}$	NNLO_{opt}	Exp.	Ref.
a_{pp}^C	-7.8188	-7.8174	-7.8196(26)	[25]
			-7.8149(29)	[26]
r_{pp}^C	2.795	2.755	2.790(14)	[25]
			2.769(14)	[26]
a_{pp}^N	-17.083	-17.825		
r_{pp}^N	2.876	2.817		
a_{nn}	-18.900	-18.889	-18.95(40)	[27, 28]
r_{nn}	2.838	2.797	2.75(11)	[29]
a_{np}	-23.732	-23.749	-23.740(20)	[23]
r_{np}	2.725	2.684	2.77(5)	[23]
B_D (MeV)	2.224575	2.224582	2.224575(9)	[23]
r_D (fm)	1.975	1.967	1.97535(85)	[30]
Q_D (fm^2)	0.275	0.272	0.2859(3)	[23]
P_D (%)	4.51	4.05		

Figure 1 shows some np phase shifts of NNLO_{opt} and compares them to phase shifts from other potentials and partial wave analyses. Apart from the 3P -waves, the phase shifts of NNLO_{opt} are in very close agreement with the ones obtained at N^3LO . Note, however, that these deviations do not spoil the good χ^2 at laboratory energies below 125 MeV.




FIG. 1. (Color online) Computed np phase shifts of the optimized NNLO potential of this work (red), the NNLO potential of Ref. [3] (dashed, blue), and the N^3LO potential [4] (green, dotted) compared to the Nijmegen phase shift analysis [18] (solid dots) and the VPI/GWU analysis SM99 [31] (open circles).

TABLE IV. Ground-state energies (in MeV) and point proton radii (in fm) for ${}^3\text{H}$, ${}^3\text{He}$, and ${}^4\text{He}$ using the NNLO_{opt} with and without the NNLO 3NF interaction for $c_D = -0.20$ and $c_E = -0.36$.

	$E({}^3\text{H})$	$E({}^3\text{He})$	$E({}^4\text{He})$	$r_p({}^4\text{He})$
NNLO	-8.249	-7.501	-27.759	1.43(8)
NNLO+NNN	-8.469	-7.722	-28.417	1.43(8)
Experiment	-8.482	-7.717	-28.296	1.467(13)

Three-nucleon forces also appear at NNLO , and two additional LECs (c_D and c_E) enter. These are determined from calculations in the three-nucleon and four-nucleon systems. We find that the binding energies of ${}^3\text{H}$, ${}^3\text{He}$, and ${}^4\text{He}$ do not uniquely determine c_D and c_E , and the parametric dependence of both LECs is very similar to the ones found in previous studies [6, 32, 33]. Due to this close similarity, we choose $c_D = -0.2$ guided by the triton half life [33] and obtain $c_E = -0.36$ from optimization to the binding energies. The resulting point charge radii of ${}^4\text{He}$ is also in good agreement with experiment, see Table IV.

Performance of NNLO_{opt} for light and medium-mass nuclei and neutron matter – In this paper, we apply NNLO_{opt} to ${}^{10}\text{B}$, isotopes of oxygen and calcium, and neutron matter. The considered systems are particularly interesting because the current NN chiral interactions at N^3LO completely fail to describe key aspects of their structure.

To study the ground- and first excited state in ${}^{10}\text{B}$ we carry out no-core shell model (configuration interaction) calculations [34] using the bare NNLO_{opt} in model spaces of up to $N_{\text{max}} = 10$ harmonic oscillator (HO) shells ($10 \hbar\Omega$) above the unperturbed configuration. These model spaces are not large enough to provide fully converged results for the ground- and first excited state of ${}^{10}\text{B}$. Still, the variational upper bounds for the energies are -54.35 MeV for the 1^+ state and -54.32 MeV for the 3^+ state. The energies are very close, in contrast to $\text{N}^3\text{LO}_{\text{EM}}$, which yields a level spacing of about 1.2 MeV between the $J^\pi = 1^+$ ground state and the $J^\pi = 3^+$ excited state [6].

Chiral NN interactions at N^3LO fail to explain the neutron drip-line in oxygen isotopes and 3NFs have been the key element for the understanding of the structure of nuclei around ${}^{24}\text{O}$ [7, 8]. Figure 2 shows the experimental ground-state energies of oxygen isotopes, and compares the results from coupled-cluster (CC) computations in the Λ triples approximation [35–37]. Our CC calculations employ a Hartree-Fock basis (HF) built from $N_{\text{max}} = 15$ HO shells at $\hbar\Omega = 20$ MeV. Due to the “softness” of NNLO_{opt} , this model space is sufficiently large to converge the ground- and excited states of the nuclei considered. In addition, we performed shell-model (SM) calculations assuming the closed ${}^{16}\text{O}$ core with an effective interaction derived from many-body perturbation theory to third order in the interaction and including folded diagrams [38]. For the SM calculations, the single-particle energies were taken from the experimental ${}^{17}\text{O}$ spectrum. In both CC and SM, NNLO_{opt} results are close to experiment. This is in contrast to the $\text{N}^3\text{LO}_{\text{EM}}$ case, which requires 3NFs to provide reasonable description of measured values.

Now we consider the heavy isotopes of calcium. Here, ${}^{48}\text{Ca}$ is doubly magic, ${}^{52}\text{Ca}$ exhibits a soft subshell closure, and ${}^{54}\text{Ca}$ is predicted to have an even softer subshell closure [10]. A signature of shell closure is the location of the first 2^+ state. We employed CC equation-of-motion methods within the singles- and doubles approximation [37, 39] to compute the first 2^+ state in the calcium isotopes. Figure 3 shows that $\text{N}^3\text{LO}_{\text{EM}}$ fails to describe the location of the first 2^+ state in ${}^{40,48,50,52,54,56}\text{Ca}$. In contrast, NNLO_{opt} yields ${}^{48}\text{Ca}$ as a doubly magic nucleus, and predicts subshell closures in ${}^{52,54}\text{Ca}$. The NNLO_{opt} overbinds the calcium isotopes by about 1 MeV per nucleon. In particular ${}^{40,48,52}\text{Ca}$ are overbound by 1.03 MeV, 1.06 MeV and 1.04 MeV per nucleon, respectively. That is, the excess energy per nucleon is fairly con-




FIG. 2. (Color online). The ground-state energies of oxygen isotopes obtained in CC with the NNLO_{opt} and $\text{N}^3\text{LO}_{\text{EM}}$ interactions compared to experiment. The inset shows SM results.




FIG. 3. (Color online). The first 2_1^+ state in selected calcium isotopes obtained in CC with the NNLO_{opt} and $\text{N}^3\text{LO}_{\text{EM}}$ interactions compared to experiment.

stant; hence, NNLO_{opt} reproduces binding energy differences, such as neutron-separation energies and low-lying excited states rather well.

The complete description of nuclei at NNLO also requires 3NFs. We computed the first 2^+ state in $^{22,24}\text{O}$ and in ^{48}Ca with the 3NF compatible with the NNLO_{opt} interaction. The matrix elements of the 3NF are very expensive computationally, and we must presently limit their calculation to three-body energies up to $e_{3\text{max}} = 2n_a + l_a + 2n_b + l_b + 2n_c + l_c = 14$. (Recall that we employ 15 major harmonic oscillator shells for the NN interaction.) We also used the normal ordered two-body approximation for the 3NF [40, 41] with respect to a HF reference. With the restriction of $e_{3\text{max}} = 14$, we were not able to obtain fully converged results for the binding energies of oxygen and calcium isotopes. How-

ever, excitation energies relative to the ground state converge somewhat better. Our results for the first 2^+ state in $^{22,24}\text{O}$ and in ^{48}Ca are 2.3(3) MeV, 3.5(5) MeV and 4.8(7) MeV, respectively. We estimate the uncertainty by varying $\hbar\Omega$ in the interval 16 – 22 MeV. The results obtained using NNLO_{opt} NN interaction alone, yields 2.5 MeV, 5.0 MeV and 4.5 MeV in $^{22,24}\text{O}$ and ^{48}Ca , respectively. These preliminary results suggest that the 3NFs may not dramatically change the results that were obtained with the NNLO_{opt} NN interaction alone.

It is instructive to compare the predictions of NNLO_{opt} and $\text{N}^3\text{LO}_{\text{EM}}$ for the neutron matter equation of state at sub-saturation densities with the results of ab-initio calculations of Refs. [42]. Figure 4 shows that the performance of NNLO_{opt} is on par with the EGM results of Ref. [42], which take into account the effects of 3NFs and 4NFs. The predictions of $\text{N}^3\text{LO}_{\text{EM}}$ deviate from other results at higher densities.




FIG. 4. (Color online). Energy per nucleon for neutron matter for NNLO_{opt} and $\text{N}^3\text{LO}_{\text{EM}}$ [4]. The calculations used the CC method with the inclusion of particle-particle ladders and a continuous single-particle spectrum. The shaded area (EGM) show uncertainty bands for N^3LO chiral effective field theory calculations of Ref. [42], including 3NFs.

Conclusions. – We constructed the new NN chiral EFT interaction NNLO_{opt} at next-to-next-to-leading order using the optimization tool POUNDers in the phase-shift analysis. The optimization of the low-energy constants in the NN -sector at NNLO yields a χ^2/datum of about one for laboratory scattering energies below 125 MeV. The NNLO_{opt} NN interaction yields a very good agreement with binding energies and radii for $A = 3, 4$ nuclei. Key aspects of nuclear structure such as excitation spectra, the position of the neutron drip line in oxygen, shell-closures in calcium, and the neutron matter equation of state at sub-saturation densities are repro-

duced by NNLO_{opt} interaction alone, without resorting to 3NFs. We performed the initial calculation of the first 2^+ states in $^{22,24}\text{O}$ and ^{48}Ca with NNLO_{opt} supplemented by a 3NF, and found effects of 3NFs to be small and a good agreement with experimental excitation energies. The precise role of 3NFs in medium-mass nuclei, the quantification of theoretical uncertainties, and optimizations at higher-order chiral interactions, will be addressed in forthcoming investigations.

We thank M. P. Kartamyshev, B. D. Carlsson, and H. T. Johansson for discussions and related code development. This work was supported by the Research Council of Norway under contract ISP-Fysikk/216699, by the Office of Nuclear Physics, U.S. Department of Energy (Oak Ridge National Laboratory), under Grant Nos. DE-FG02-03ER41270 (University of Idaho), DE-FG02-96ER40963 (University of Tennessee), DE-AC02-06CH11357 (Argonne), and DE-SC0008499 (NUCLEI SciDAC-3 Collaboration), the Swedish Research Council (dnr 2007-4078), and by the European Research Council (ERC-StG-240603). Computer time was provided by the Innovative and Novel Computational Impact on Theory and Experiment (INCITE) program. This research used resources of the Oak Ridge Leadership Computing Facility located in the Oak Ridge National Laboratory, which is supported by the Office of Science of the Department of Energy under Contract No. DE-AC05-00OR22725 and used computational resources of the National Center for Computational Sciences, the National Institute for Computational Sciences, and the Notur project in Norway.

-
- [1] U. van Kolck, Phys. Rev. C **49**, 2932 (1994).
 [2] E. Epelbaum, H.-W. Hammer, and U.-G. Meißner, Rev. Mod. Phys. **81**, 1773 (2009).
 R. Machleidt and D. Entem, Phys. Rep. **503**, 1 (2011).
 [3] E. Epelbaum, W. Glöckle, and U.-G. Meißner, Eur. Phys. J. A **19**, 401 (2004).
 E. Epelbaum, A. Nogga, W. Glöckle, H. Kamada, U.-G. Meißner, and H. Witała, Eur. Phys. J. A **15**, 543 (2002), ISSN 1434-6001.
 [4] D. R. Entem and R. Machleidt, Phys. Rev. C **68**, 041001(R) (2003).
 [5] H.-W. Hammer, A. Nogga, and A. Schwenk, Rev. Mod. Phys. **85**, 197 (2013).
 [6] P. Navrátil, V. G. Gueorguiev, J. P. Vary, W. E. Ormand, and A. Nogga, Phys. Rev. Lett. **99**, 042501 (2007).
 [7] T. Otsuka, T. Suzuki, J. D. Holt, A. Schwenk, and Y. Akaishi, Phys. Rev. Lett. **105**, 032501 (2010).
 [8] G. Hagen, M. Hjorth-Jensen, G. R. Jansen, R. Machleidt, and T. Papenbrock, Phys. Rev. Lett. **108**, 242501 (2012).
 [9] J. D. Holt, T. Otsuka, A. Schwenk, and T. Suzuki, J. Phys. G **39**, 085111 (2012).
 [10] G. Hagen, M. Hjorth-Jensen, G. R. Jansen, R. Machleidt, and T. Papenbrock, Phys. Rev. Lett. **109**, 032502 (2012).
 [11] S. Ishikawa and M. R. Robilotta, Phys. Rev. C **76**, 014006 (2007).
 [12] V. Bernard, E. Epelbaum, H. Krebs, and U.-G. Meißner, Phys. Rev. C **77**, 064004 (2008).
 V. Bernard, E. Epelbaum, H. Krebs, and U.-G. Meißner, Phys. Rev. C **84**, 054001 (2011).
 [13] H. Krebs, A. Gasparyan, and E. Epelbaum, Phys. Rev. C **85**, 054006 (2012).
 [14] E. Epelbaum, A. Nogga, W. Glöckle, H. Kamada, U.-G. Meißner, and H. Witała, Phys. Rev. C **66**, 064001 (2002).
 [15] M. Kortelainen, T. Lesinski, J. Moré, W. Nazarewicz, J. Sarich, N. Schunck, M. V. Stoitsov, and S. Wild, Phys. Rev. C **82**, 024313 (2010).
 [16] P.-G. Reinhard and W. Nazarewicz, Phys. Rev. C **81**, 051303 (2010).
 [17] T. Munson, J. Sarich, S. M. Wild, S. Benson, and L. Curfman McInnes, Technical Memorandum ANL/MCS-TM-322, Argonne National Laboratory, Argonne, Illinois (2012), see <http://www.mcs.anl.gov/tao>.
 [18] V. G. J. Stoks, R. A. M. Klomp, M. C. M. Rentmeester, and J. J. de Swart, Phys. Rev. C **48**, 792 (1993).
 [19] C. M. Vincent and S. C. Phatak, Phys. Rev. C **10**, 391 (1974).
 [20] D. H. Lu, T. Mefford, R. H. Landau, and G. Song, Phys. Rev. C **50**, 3037 (1994).
 [21] P. Büttiker and U.-G. Meißner, Nuclear Physics A **668**, 97 (2000).
 [22] A. Ekström et al., to be published (2013).
 [23] R. Machleidt, Phys. Rev. C **63**, 024001 (2001).
 [24] G. Cox, G. Eaton, C. V. Zyl, O. Jarvis, and B. Rose, Nuclear Physics B **4**, 353 (1967), 21 pp diff. cross section data at 144.1 MeV.
 O. Jarvis, C. Whitehead, and M. Shah, Physics Letters B **36**, 409 (1971), 26 pp diff. cross section data at 144.0 MeV.
 [25] J. R. Bergervoet, P. C. van Campen, W. A. van der Sanden, and J. J. de Swart, Phys. Rev. C **38**, 15 (1988).
 [26] W. A. van der Sanden, A. H. Emmen, and J. J. de Swart, Tech. Rep., Nijmegen (1983), unpublished.
 [27] D. E. Gonzalez Trotter, F. S. Meneses, W. Tornow, C. R. Howell, Q. Chen, A. S. Crowell, C. D. Roper, R. L. Walter, D. Schmidt, H. Witała, et al., Phys. Rev. C **73**, 034001 (2006).
 [28] Q. Chen, C. R. Howell, T. S. Carman, W. R. Gibbs, B. F. Gibson, A. Hussein, M. R. Kiser, G. Mertens, C. F. Moore, C. Morris, et al., Phys. Rev. C **77**, 054002 (2008).
 [29] G. Miller, B. Nefkens, and I. Šlaus, Phys. Rep. **194**, 1 (1990).
 [30] A. Huber, T. Udem, B. Gross, J. Reichert, M. Kourogi, K. Pachucki, M. Weitz, and T. W. Hänsch, Phys. Rev. Lett. **80**, 468 (1998).
 [31] R. A. Arndt, I. I. Strakovsky, and R. L. Workman (1999), SAID, Scattering Analysis Interactive Dial-in computer facility, George Washington University (formerly Virginia Polytechnic Institute), solution SM99 (Summer 1999); for more information see, e. g., R. A. Arndt, I. I. Strakovsky, and R. L. Workman, Phys. Rev. C **50**, 2731 (1994).
 [32] A. Nogga, P. Navrátil, B. R. Barrett, and J. P. Vary, Phys. Rev. C **73**, 064002 (2006).
 [33] D. Gazit, S. Quaglioni, and P. Navrátil, Phys. Rev. Lett. **103**, 102502 (2009).
 [34] B. R. Barrett, P. Navrátil, and J. P. Vary, Prog. Part. Nucl. Phys. **69**, 131 (2013).
 [35] S. A. Kucharski and R. J. Bartlett, The J. Chem. Phys. **108**, 5243 (1998).

- [36] A. G. Taube and R. J. Bartlett, *The J. Chem. Phys.* **128**, 044110 (2008).
- [37] G. Hagen, T. Papenbrock, D. J. Dean, and M. Hjorth-Jensen, *Phys. Rev. C* **82**, 034330 (2010).
- [38] M. Hjorth-Jensen, T. T. S. Kuo, and E. Osnes, *Phys. Rep.* **261**, 125 (1995).
- [39] G. R. Jansen, M. Hjorth-Jensen, G. Hagen, and T. Papenbrock, *Phys. Rev. C* **83**, 054306 (2011).
- [40] G. Hagen, T. Papenbrock, D. J. Dean, A. Schwenk, A. Nogga, M. Włoch, and P. Piecuch, *Phys. Rev. C* **76**, 034302 (2007).
- [41] S. Binder, J. Langhammer, A. Calci, P. Navrátil, and R. Roth, *Phys. Rev. C* **87**, 021303 (2013).
- [42] I. Tews, T. Krüger, K. Hebeler, and A. Schwenk, *Phys. Rev. Lett.* **110**, 032504 (2013).

The submitted manuscript has been created by the University of Chicago as Operator of Argonne National Laboratory ("Argonne") under Contract DE-AC02-06CH11357 with the U.S. Department of Energy. The U.S. Government retains for itself, and others acting on its behalf, a paid-up, nonexclusive, irrevocable worldwide license in said article to reproduce, prepare derivative works, distribute copies to the public, and perform publicly and display publicly, by or on behalf of the Government.